

Table 3.72

Estimated prevalence and most recent use of cocaine and crack

By sex, race, ethnicity, age, and region, United States, 1995

	Cocaine ^a				Crack			
	Never used	Ever used	Most recent use		Never used	Ever used	Most recent use	
			Within last year	Within last 30 days			Within last year	Within last 30 days
Total (N=17,747)	89.7%	10.3%	1.7%	0.7%	98.2%	1.8%	0.5%	0.2%
Sex								
Male	87.2	12.8	2.3	1.0	97.4	2.6	0.6	0.3
Female	92.1	7.9	1.2	0.4	98.9	1.1	0.4	0.1
Race, ethnicity								
White	88.7	11.3	1.7	0.6	98.3	1.7	0.4	0.1
Black	91.9	8.1	1.9	1.1	97.0	3.0	1.0	0.6
Hispanic	92.5	7.5	1.8	0.7	98.3	1.7	0.5	0.2
Age								
12 to 17 years	98.0	2.0	1.7	0.8	99.1	0.9	0.6	0.2
18 to 25 years	90.2	9.8	4.3	1.3	97.1	2.9	1.1	0.3
26 to 34 years	78.4	21.6	3.1	1.2	95.8	4.2	0.9	0.3
35 years and older	91.4	8.6	0.8	0.4	98.9	1.1	0.2	0.2
Region								
Northeast	88.8	11.2	1.5	0.5	98.2	1.8	0.4	0.1
North Central	91.1	8.9	1.6	0.7	98.7	1.3	0.4	0.1
South	91.7	8.3	1.5	0.8	98.0	2.0	0.6	0.3
West	86.1	13.9	2.5	0.7	97.8	2.2	0.5	0.2

Note: See Note, table 3.69. For survey methodology, see Appendix 8.

^aThe estimates for cocaine include crack.

Source: U.S. Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, *National Household Survey on Drug Abuse: Population Estimates 1995* (Rockville, MD: U.S. Department of Health and Human Services, 1996), pp. 29-39. Table constructed by SOURCEBOOK staff.